

D-branes in a plane-wave background

M.R. Gaberdiel^{1 §}, M.B. Green²

¹ Department of Mathematics
King's College, London

Strand, London, WC2R 2LS, UK

² Department of Applied Mathematics and Theoretical Physics
Cambridge University

Wilberforce Road, Cambridge, CB3 0WA, UK

Abstract: The D-branes of the maximally supersymmetric plane-wave background are described.

1 Introduction

The study of type IIB superstring theory in plane wave backgrounds is a fertile area for investigating the properties of string theory with nontrivial Ramond–Ramond (RR) condensates. One of the best studied examples is the Penrose limit of $AdS_5 \times S^5$ [1], for which the dual CFT has a corresponding limit [2]. In the original $AdS_5 \times S^5$ background this correspondence is difficult to check quantitatively since string theory in the $AdS_5 \times S^5$ background has not, so far, proved tractable. On the other hand, the plane-wave theory can be formulated as a free two-dimensional field theory, at least in the light-cone gauge, and is therefore exactly solvable [3, 4].

In this talk we shall describe the construction of supersymmetric D-branes for this string theory background. Since the world-sheet theory in light cone gauge is not conformally invariant, a number of modifications arise relative to the usual conformal field theory discussion. In particular, the functions that enter in the calculation of the cylinder amplitudes are not standard θ functions, and the open-closed consistency condition follows from rather non-trivial modular transformation properties of deformed θ functions.

This talk is based on [5] and [6].

2 Notation and review

The pp -wave background is a ten-dimensional space-time with metric,

$$ds^2 = 2dx^+ dx^- - \mu^2 x^I x^I dx^+ dx^+ + dx^I dx^I, \quad (1)$$

where $x^\pm = (x^0 \pm x^9)/\sqrt{2}$ and $I = 1, \dots, 8$. The five-form RR field strength is given by

$$F_{+1234} = F_{+5678} = 2\mu, \quad (2)$$

[§]corresponding author : mrg@mth.kcl.ac.uk

where μ is a constant. In light-cone gauge where $x^+ = 2\pi p^+ \tau$, the Lagrangian describes eight massive free scalar and eight massive free fermion fields

$$\mathcal{L} = \frac{1}{4\pi} \left(\partial_+ x^I \partial_- x^I - m^2 (x^I)^2 \right) + \frac{i}{2\pi} \left(S^a \partial_+ S^a + \tilde{S}^a \partial_- \tilde{S}^a - 2m S^a \Pi_{ab} \tilde{S}^b \right), \quad (3)$$

where S^a and $\tilde{S}^a$ are $SO(8)$ spinors of the same chirality and $\Pi = \gamma^1 \gamma^2 \gamma^3 \gamma^4$. The mass parameter m is defined by $m = 2\pi p^+ \mu$. The 8×8 matrices, γ_{ab}^I and $\gamma_{\dot{a}\dot{b}}^I$, are the off-diagonal blocks of the 16×16 $SO(8)$ γ -matrices and couple $SO(8)$ spinors of opposite chirality. The presence of Π in the fermionic sector of the lagrangian breaks the symmetry from $SO(8)$ to $SO(4) \times SO(4)$.

The modes of the transverse bosonic coordinates, x^I , of a string in type IIB pp -wave light-cone gauge string theory [3] are α_k^I and $\tilde{\alpha}_k^I$, where I is a vector index of $SO(8)$, and $k \in \mathbb{Z}$ with $k \neq 0$. These modes satisfy the commutation relations

$$[\alpha_k^I, \alpha_l^J] = \omega_k \delta^{IJ} \delta_{k,-l}, \quad [\alpha_k^I, \tilde{\alpha}_l^J] = 0, \quad [\tilde{\alpha}_k^I, \tilde{\alpha}_l^J] = \omega_k \delta^{IJ} \delta_{k,-l}, \quad (4)$$

where

$$\omega_k = \text{sign}(k) \sqrt{k^2 + m^2} \quad |k| > 0. \quad (5)$$

In addition there are the bosonic zero modes that describe the centre of mass position x_0^I and some generalised momentum p_0^I with $[p_0^I, x_0^J] = -i\delta^{IJ}$. It is convenient to introduce the creation and annihilation operators

$$a_0^I = \frac{1}{\sqrt{2m}}(p_0^I + imx_0^I), \quad \bar{a}_0^I = \frac{1}{\sqrt{2m}}(p_0^I - imx_0^I), \quad (6)$$

in terms of which the commutation relation is then $[\bar{a}_0^I, a_0^J] = \delta^{IJ}$. The modes of the fermionic fields S_k^a and $\tilde{S}_k^a$, where a is a spinor index of $SO(8)$ and $k \in \mathbb{Z}$, satisfy the anti-commutation relations

$$\{S_k^a, S_l^b\} = \delta^{ab} \delta_{k,-l}, \quad \{S_k^a, \tilde{S}_l^b\} = 0, \quad \{\tilde{S}_k^a, \tilde{S}_l^b\} = \delta^{ab} \delta_{k,-l}. \quad (7)$$

It is convenient to introduce the zero-mode combinations $\theta_0^a = \frac{1}{\sqrt{2}}(S_0^a + i\tilde{S}_0^a)$ as well as $\bar{\theta}_0^a = \frac{1}{\sqrt{2}}(S_0^a - i\tilde{S}_0^a)$, and further

$$\begin{aligned} \theta_R &= \frac{1}{2}(1 + \Pi)\theta_0, & \bar{\theta}_R &= \frac{1}{2}(1 + \Pi)\bar{\theta}_0, \\ \theta_L &= \frac{1}{2}(1 - \Pi)\theta_0, & \bar{\theta}_L &= \frac{1}{2}(1 - \Pi)\bar{\theta}_0. \end{aligned} \quad (8)$$

In light-cone gauge the thirty-two components of the supersymmetries decompose into ‘dynamical’ and ‘kinematical’ components. The ‘dynamical’ supercharges, $Q_{\dot{a}}$ and $\tilde{Q}_{\dot{a}}$ ($\dot{a} = 1, \dots, 8$), anticommute to give the light-cone hamiltonian

$$\begin{aligned} 2p^+ H &= m \left(a_0^I \bar{a}_0^I + i S_0^a \Pi_{ab} \tilde{S}_0^b + 4 \right) + \sum_{k=1}^{\infty} \left[\alpha_{-k}^I \alpha_k^I + \tilde{\alpha}_{-k}^I \tilde{\alpha}_k^I + \omega_k \left(S_{-k}^a S_k^a + \tilde{S}_{-k}^a \tilde{S}_k^a \right) \right] \\ &= m \left(a_0^I \bar{a}_0^I + \theta_L^a \bar{\theta}_L^a + \bar{\theta}_R^a \theta_R^a \right) + \sum_{k=1}^{\infty} \left[\alpha_{-k}^I \alpha_k^I + \tilde{\alpha}_{-k}^I \tilde{\alpha}_k^I + \omega_k \left(S_{-k}^a S_k^a + \tilde{S}_{-k}^a \tilde{S}_k^a \right) \right]. \end{aligned} \quad (9)$$

In addition the theory has sixteen ‘kinematical’ supercharges $Q_a \equiv S_0^a$ and $\tilde{Q}_a \equiv \tilde{S}_0^a$, where $a = 1, \dots, 8$. The dynamical supercharges commute with the light-cone hamiltonian, but the kinematical supercharges do not.

3 The construction of the boundary states

D-branes in string theory can always be described by boundary states. This description makes the coupling of the D-brane to the closed string states of the theory manifest. In the following we shall only be discussing instantonic D-branes, which are defined by the embeddings of euclidean $(p+1)$ -dimensional world-volumes.¹ These are the cases in which the light-cone directions $x^\pm$ are orthogonal to the brane. We will adopt a notation, following [7], in which these instantonic branes are denoted (r, s) -branes ($r+s=p+1$), where r and s are the numbers of Neumann directions associated with the two $SO(4)$ factors in the transverse space.

The boundary states are (up to important normalisations) uniquely determined by the gluing conditions that describe how left- and right-moving fields are related at the boundary. Here we shall only consider D-branes that preserve half of the dynamical supersymmetries, *i.e.*

$$\left(Q_{\dot{a}} + i\eta M_{\dot{a}\dot{b}} \tilde{Q}_{\dot{b}} \right) \|(r, s), \eta \rangle = 0, \quad (10)$$

where the value of $\eta = \pm 1$ distinguishes a brane from an anti-brane. The matrix M is given by

$$M_{\dot{a}\dot{b}} = \left(\prod_{I \in \mathcal{N}} \gamma^I \right)_{\dot{a}\dot{b}}, \quad (11)$$

where the product extends over the Neumann directions $\mathcal{N}$. In order for the boundary state to define a standard D-brane one requires in addition that the bosonic degrees of freedom satisfy

$$\begin{aligned} \left(\alpha_k^I - \tilde{\alpha}_{-k}^I \right) \|(r, s), \mathbf{y}_t \rangle &= 0, \quad k \in \mathbb{Z}, \\ \left(\bar{a}_0^I - a_0^I + i\sqrt{2m}y_t^I \right) \|(r, s), \mathbf{y}_t \rangle &\equiv -i\sqrt{2m}(x_0^I - y_t^I) \|(r, s), \mathbf{y}_t \rangle = 0, \end{aligned} \quad (12)$$

for each Dirichlet direction I , as well as

$$\begin{aligned} \left(\alpha_k^J + \tilde{\alpha}_{-k}^J \right) \|(r, s), \mathbf{y}_t \rangle &= 0, \quad k \in \mathbb{Z}, \\ \left(\bar{a}_0^J + a_0^J \right) \|(r, s), \mathbf{y}_t \rangle &= 0 \end{aligned} \quad (13)$$

for each Neumann direction J . (For simplicity we are considering here the case without Wilson lines.) It is easy to see that (10), together with (12) and (13) implies that

$$\left(S_0^a + i\eta M_{ab} \tilde{S}_0^b \right) \|(r, s), \mathbf{y}_t \rangle = 0, \quad (14)$$

where M_{ab} is the same product of γ matrices as in (11). This last equation implies that a complex combination of the kinematical supersymmetries is preserved by the boundary state. The structure of the boundary states depends now crucially on the choice of M .

Class I: The first class is the one that was studied in [8, 9, 5] and arises when the matrix M_{ab} satisfies

$$\Pi M \Pi M = -1, \quad (15)$$

where $\Pi = \gamma^1 \gamma^2 \gamma^3 \gamma^4$. From a supergravity point of view, these branes were analysed in [7]. The condition (15) is equivalent to $|r-s|=2$. The branes of this kind satisfy the standard fermionic gluing condition

$$\left(S_n^a + i\eta M_{ab} \tilde{S}_{-n}^b \right) \|(r, s), \mathbf{y}_t \rangle = 0. \quad (16)$$

¹As is explained in [5, 6], the following discussion can be easily generalised for time-like branes.

They satisfy (10) if $y_t^I = 0$ for the Dirichlet directions, and the corresponding open string preserves eight components (*i.e.* half) of both the dynamical and kinematical supersymmetries. A characteristic feature of this class is that the kinematical conditions (14) are not preserved as a function of x^+ since the commutator with the light-cone hamiltonian has the form

$$[H, Q_a + i\eta M_{ab} \tilde{Q}_b] = \frac{m\eta}{2p^+} (\Pi M^t)_{ab} (Q_b - i\eta M_{bc} \tilde{Q}_c) . \quad (17)$$

In this case the open-string theory has a mass term in its hamiltonian of the form $S_0 M \Pi S_0$ [9], and the ground state is an unmatched boson. In particular, the open string one-loop amplitude (and the corresponding closed string cylinder diagram) therefore does not vanish.

Class II: The second class arises when the matrix M_{ab} satisfies

$$\Pi M \Pi M = 1 , \quad (18)$$

a possibility that was not considered in [8, 9, 5] but arose in the supergravity analyses of [7, 10] and was later analysed in detail in [6] (the open string description was independently worked out in [11]). In this case $|r-s| = 0, 4$. The only branes of this type that satisfy (10) are the $(0, 0)$ brane at an arbitrary position², for which the fermionic gluing conditions are

$$(S_n^a + i\eta R_n^{ab} \tilde{S}_{-n}^b) \|(0, 0), \mathbf{y}\rangle\rangle = 0 , \quad (19)$$

and R_n is the matrix

$$R_n = \frac{1}{n} (\omega_n \mathbb{1} - \eta m \Pi) . \quad (20)$$

In addition, the $(4, 0)$ and $(0, 4)$ branes satisfy (10) provided that the Neumann boundary conditions (13) are suitably modified (see [6]). The corresponding open strings preserve half of the dynamical supersymmetries, but none of the kinematical supersymmetries. However, in contradistinction to class I,

$$[H, Q_a + i\eta M_{ab} \tilde{Q}_b] = -\frac{m\eta}{2p^+} (\Pi M^t)_{ab} (Q_b + i\eta M_{bc} \tilde{Q}_c) , \quad (21)$$

and thus the kinematical conditions (14) are preserved as a function of x^+ . In this case $S_0 M \Pi S_0 \equiv 0$ and the open-string mass term vanishes, thus giving true fermionic zero modes (see also [12]). The ground states then form a degenerate supermultiplet, and the one-loop open string amplitude (as well as the corresponding closed string cylinder diagram) vanishes.

4 Open-closed duality

One of the most important consistency conditions for D-branes is the so-called open-closed duality relation. It requires that the cylinder diagram, that describes the interaction energy between two D-branes, can be evaluated in two different ways. From the closed string point of view, the cylinder diagram is the tree-level diagram describing the exchange of closed string states between two external boundary states. On the other hand, the diagram can also be interpreted as a one-loop open string diagram. These two points of view are related by exchanging the roles of the world-sheet parameters τ and σ . This

²The same also holds for the $(4, 4)$ brane, but the corresponding boundary state does not have a sensible flat space limit and is therefore probably inconsistent.

transformation replaces the modular parameter t of the cylinder by $\tilde{t} = 1/t$; it also replaces the mass-parameter m by $\widehat{m} = mt$. (See [5, 6] for more details.)

In the usual flat space background, the open-closed duality condition is satisfied since the cylinder amplitude can be written in terms of standard θ functions that have well understood transformation properties under the modular transformation $t \mapsto \tilde{t}$. For the case of the plane-wave background, on the other hand, the cylinder amplitudes involve non-trivial deformations of θ functions. In the case of the class I branes³, the relevant functions are

$$f_1^{(m)}(q) = q^{-\Delta_m} (1 - q^m)^{\frac{1}{2}} \prod_{n=1}^{\infty} (1 - q^{\sqrt{m^2+n^2}}), \quad (22)$$

$$f_2^{(m)}(q) = q^{-\Delta_m} (1 + q^m)^{\frac{1}{2}} \prod_{n=1}^{\infty} (1 + q^{\sqrt{m^2+n^2}}), \quad (23)$$

$$f_3^{(m)}(q) = q^{-\Delta'_m} \prod_{n=1}^{\infty} \left(1 + q^{\sqrt{m^2+(n-1/2)^2}}\right), \quad (24)$$

$$f_4^{(m)}(q) = q^{-\Delta'_m} \prod_{n=1}^{\infty} \left(1 - q^{\sqrt{m^2+(n-1/2)^2}}\right), \quad (25)$$

where $q = e^{-2\pi t}$, and Δ_m and Δ'_m are given as

$$\begin{aligned} \Delta_m &= -\frac{1}{(2\pi)^2} \sum_{p=1}^{\infty} \int_0^{\infty} ds e^{-p^2 s} e^{-\pi^2 m^2/s}, \\ \Delta'_m &= -\frac{1}{(2\pi)^2} \sum_{p=1}^{\infty} (-1)^p \int_0^{\infty} ds e^{-p^2 s} e^{-\pi^2 m^2/s}. \end{aligned} \quad (26)$$

The quantities Δ_m and Δ'_m are the Casimir energies of a single (two-dimensional) boson of mass m on a cylindrical world-sheet with periodic and anti-periodic boundary conditions, respectively. For $m = 0$, Δ_m and Δ'_m simplify to the usual flat-space values,

$$\begin{aligned} \Delta_0 &= -\frac{1}{(2\pi)^2} \sum_{p=1}^{\infty} \frac{1}{p^2} = -\frac{1}{24}, \\ \Delta'_0 &= -\frac{1}{(2\pi)^2} \sum_{p=1}^{\infty} \frac{(-1)^p}{p^2} = \frac{1}{48}. \end{aligned} \quad (27)$$

Thus $f_2^{(m)}(q)$, $f_3^{(m)}(q)$ and $f_4^{(m)}(q)$ simply reduce to the standard $f_2(q)$, $f_3(q)$ and $f_4(q)$ functions [13] as $m \rightarrow 0$.

For the case of the class I branes the open-closed consistency condition is then a consequence of the remarkable transformation properties of these functions [5]

$$f_1^{(m)}(q) = f_1^{(\widehat{m})}(\tilde{q}), \quad f_2^{(m)}(q) = f_4^{(\widehat{m})}(\tilde{q}), \quad f_3^{(m)}(q) = f_3^{(\widehat{m})}(\tilde{q}), \quad (28)$$

where $\tilde{q} = e^{-2\pi\tilde{t}} = e^{-2\pi/t}$ and $\widehat{m} = mt$. In the limit $m \rightarrow 0$ the second and third equations in (28) reproduce standard θ -function identities. The identity for f_1 (or η) can also be derived from the first equation of (28). In fact, both sides of the first equation tend to zero as $m \rightarrow 0$ since $(1 - q^m)^{\frac{1}{2}} = \sqrt{2\pi t m} + \mathcal{O}(m)$ and $(1 - \tilde{q}^{\widehat{m}})^{\frac{1}{2}} = \sqrt{2\pi m} + \mathcal{O}(m)$. Thus, after dividing the first equation by $\sqrt{m}$, the limit $m \rightarrow 0$ becomes

$$f_1(q) = t^{-\frac{1}{2}} f_1(\tilde{q}), \quad (29)$$

³The functions that arise for the class II branes are yet more complicated; they are described in detail in [6], where their modular properties are also discussed.

thus reproducing the standard modular transformation property of the f_1 (or η) function.

Acknowledgement MRG would like to thank the organisers of the symposium for organising a very pleasant conference, and for giving him the opportunity to speak. He is grateful to the Royal Society for a University Research Fellowship. We also acknowledge partial support from the PPARC Special Programme Grant ‘String Theory and Realistic Field Theory’, PPA/G/S/1998/0061 and the EEC contract HPRN-2000-00122.

References

- [1] M. Blau, J. Figueroa-O’Farrill, C. Hull, G. Papadopoulos, *Penrose limits and maximal supersymmetry*, Class. Quant. Grav. **19**, L87 (2002); [hep-th/0201081](#).
- [2] D. Berenstein, J. Maldacena, H. Nastase, *Strings in flat space and pp waves from $\mathcal{N} = 4$ Super Yang Mills*, JHEP **0204**, 013 (2002); [hep-th/0202021](#).
- [3] R.R. Metsaev, *Type IIB Green-Schwarz superstring in plane wave Ramond-Ramond background*, Nucl. Phys. **B625**, 70 (2002); [hep-th/0112044](#).
- [4] R.R. Metsaev, A.A. Tseytlin, *Exactly solvable model of superstring in plane wave Ramond-Ramond background*, Phys. Rev. **D65**, 126004 (2002); [hep-th/0202109](#).
- [5] O. Bergman, M.R. Gaberdiel, M.B. Green, *D-brane interactions in type IIB plane-wave background*, [hep-th/0205183](#).
- [6] M.R. Gaberdiel, M.B. Green, *The D-instanton and other supersymmetric D-branes in IIB plane-wave string theory*, [hep-th/0211122](#).
- [7] K. Skenderis, M. Taylor, *Branes in AdS and pp-wave spacetimes*, JHEP **0206**, 025 (2002); [hep-th/0204054](#).
- [8] M. Billo, I. Pesando, *Boundary states for GS superstrings in an Hpp wave background*, Phys. Lett. **B536**, 121 (2002); [hep-th/0203028](#).
- [9] A. Dabholkar, S. Parvizi, *Dp branes in PP-wave background*, [hep-th/0203231](#).
- [10] P. Bain, P. Meessen, M. Zamaklar, *Supergravity solutions for D-branes in Hpp-wave backgrounds*, [hep-th/0205106](#).
- [11] K. Skenderis, M. Taylor, *Open strings in the plane wave background I: Quantization and symmetries*, [hep-th/0211011](#).
- [12] P. Bain, K. Peeters, M. Zamaklar, *D-branes in a plane wave from covariant open strings*, [hep-th/0208038](#).
- [13] J. Polchinski, Y. Cai, *Consistency of open superstring theories*, Nucl. Phys. **B296**, 91 (1988).